

663 Anderson Ferry Rd. Cincinnati, OH 45238 7074 Harrison Ave. Cincinnati, OH 45247 Phone: 513-922-8200 Fax: 513-347-2407

## **Initial Teacher Vanderbilt Assessment Scale**

Patient's Name:	Date of Birth:	Today's Dat	e:
Teacher's Name:	School:	Subject:	Class Time:

**Directions:** Each rating should be considered in the context of what is appropriate for the age of the child you are rating and should reflect that child's behavior since the beginning of the school year. Please indicate the number of weeks or months you have been able to evaluate the behaviors: \_\_\_\_\_\_

Is this evaluation based on a time the child \_\_\_\_\_ was on medication \_\_\_\_\_ was not on medication \_\_\_\_\_ not sure

Symptoms	Never	Occasionally	Often	Very Often
1. Fails to give attention to details or makes careless mistakes in schoolwork.	0	1	2	3
2. Has difficulty keeping attention to what needs to be done.	0	1	2	3
3. Does not seem to listen when spoken to directly.	0	1	2	3
4. Does not follow through when given directions and fails to finish schoolwork. (not due to oppositional behavior or failure to understand).	0	1	2	3
5. Has difficulty organizing tasks and activities.	0	1	2	3
6. Avoids, dislikes, or is reluctant to engage in tasks that require sustained mental effort.	0	1	2	3
<ol> <li>Loses things necessary for tasks or activities (e.g., school assignments, pencils, or books).</li> </ol>	0	1	2	3
8. Is easily distracted by extraneous stimuli.	0	1	2	3
9. Is forgetful in daily activities.	0	1	2	3
	-		-	
10. Fidgets with hands or feet or squirms in seat.	0	1	2	3
11. Leaves seat in classroom or in other situations in which remaining seated is expected.	0	1	2	3
12. Runs about or climbs too much when remaining seated is expected.	0	1	2	3
13. Has difficulty playing or engaging in leisure activities quietly.	0	1	2	3
14. Is "on the go" or often acts as if "driven by a motor."	0	1	2	3
15. Talks excessively.	0	1	2	3
16. Blurts out answers before questions have been completed.	0	1	2	3
17. Has difficulty waiting in line.	0	1	2	3
18. Interrupts or intrudes in others conversation and/or activities.	0	1	2	3
19. Loses temper.	0	1	2	3
20. Actively defies or refuses to go along with adult requests or rules.	0	1	2	3
21. Is angry or resentful.	0	1	2	3
22. Is spiteful and vindictive.	0	1	2	3
23. Bullies, threatens, or intimidates others.	0	1	2	3
24. Initiates physical fights.	0	1	2	3
	0	1	2	3
25. Lies to obtain goods for favors or to avoid obligations (ie, "cons" others)			2	3
<ul><li>25. Lies to obtain goods for favors or to avoid obligations (ie, "cons" others)</li><li>26. Is physically cruel to people</li></ul>	0	1	2	5
	0	1	2	3

Patient's Name: \_\_\_\_

School

Date of Birth: \_\_\_\_\_ Today's Date: \_\_\_

Grade

4

4

4

4

4

Adapted from the Vanderbilt Rating Scales developed by Mark L. Wolraich, MD

5

5

5

5

5

Teacher's Name:	
-----------------	--

	school:			G	rade:		
29. Is fearful, anxious, or worried.				0	1	2	3
30. Is self-conscious or easily embarrassed.				0	1	2	3
31. Is afraid to try new things for fear of making mistakes.				0	1	2	3
32. Feels worthless or inferior.				0	1	2	3
33. Blames self for problems, feels guilty.				0	1	2	3
34. Feels lonely, unwanted, or unloved; complains that "no one loves him or her."				0	1	2	3
35. Is sad, unhappy, or depressed.				0	1	2	3
Impairment	Excellent	Above Average	Average		mewhat of Problem	Proble	matic
36. Reading	1	2	3		4	5	
37. Mathematics	1	2	3		4	5	

1

1

1

1

1

2

2

2

2

2

3

3

3

3

3

**PRE-EXISTING PROBLEMS** 

39. Relationship with peers

42. Assignment completion

40. Following directions

43. Organizational skills

41. Disrupting class

## Use the following to assess severity:

**None:** The symptom is not present.

Mild: The symptom is present but is not significant enough to cause concern to the patient, to his/her friends, or adults.

Moderate: The symptom causes some impairment of functioning or social embarrassment.

Severe: The symptom causes impairment of functioning or social embarrassment to such a degree that it requires specific treatment.

	None	Mild	Moderate	Severe
Motor Tics- repetitive movements: jerking or twitching (e.g. eye blinking, eye opening, facial or mouth twitching, shoulder or arm movements)-describe:				
Buccal–lingual movements: Tongue thrusts, jaw clenching, chewing movement besides lip/cheek biting– describe:				
Picking at skin or fingers, nail biting, lip or cheek chewing – describe:				
Worried/Anxious				
Dull, tired, listless				
Headaches				
Stomachache				
Crabby, Irritable				
Tearful, Sad, Depressed				
Socially withdrawn – decreased interaction with others				
Hallucinations (see or hear things that aren't there)				
Loss of appetite				
Trouble sleeping (time went to sleep)				

Adapted from the Pittsburgh Side-Effects Rating Scale